



Pharmacy

Opioids for Acute Pain Relief

Introduction

Opioids are 'morphine-type' medications, used to relieve acute, severe pain. You will have been given either:

- **Morphine Sulphate solution 10mg in 5mL, or**
- **Oxycodone liquid 5mg in 5mL**

Opioids should be used for a short period after surgery or serious injury to relieve acute pain. Opioids are used with other painkillers which work together to give better pain relief. They reduce pain levels to a point where you can mobilise and undertake activities of daily living. It is unlikely that they will get rid of your pain completely. Severe pain should not last long and normally gets a little better each day.

Advice

- As your body begins to heal a little each day, you should take less opioid medicine.
- **Do not take for longer than a week. If you continue to have severe pain you should discuss with your GP.**
- You may have also been given (or advised to take) paracetamol, which should be taken regularly and supplemented with the opioid medicine as needed, to help manage your pain better.
- You may have also been given (or advised to take) a non-steroidal anti-inflammatory medicine eg ibuprofen or naproxen, which should be taken regularly and supplemented with the opioid medicine as needed, to help manage your pain better.
- Continue to take your regular medication as prescribed.
- Do not start taking additional painkillers that contain codeine, dihydrocodeine or tramadol, unless advised by your doctor or you were taking these before your hospital admission. You will be more likely to get side effects.
- Do not start taking additional medicines to help you sleep, such as benzodiazepines (eg diazepam, lorazepam, temazepam, zopiclone), unless advised by your doctor or you were taking these before your hospital admission. These medicines increase the risk of drowsiness or sedation.

How should I take my opioid medicine?

You should only take liquid opioid medicines for severe pain. The opioid liquid works quickly - it should take 15-30minutes to get pain relief which then lasts around 4 hours. Always read the directions on the label to see how much to take and how often.

Always use the oral purple syringe provided to measure your dose (after use, wash the syringe with warm soapy water, rinse and leave to dry). Do not take a larger dose or take more frequently than advised on the label.

Are there any side effects?

- **Sickness (nausea)** - Try taking your medicine with or just after a snack or meal.
- **Constipation** - Make sure you are drinking plenty of water and try to include more high fibre food in your diet unless otherwise advised by the hospital. You may need a laxative if constipation is a problem.
- **Feeling sleepy or dizzy** - This is common. If affected, do not attempt to complete complex tasks, such as operating machinery, or even boiling a kettle. **Do NOT drive.** Drowsiness (or sedation) can be an indication that you have taken too much opioid. Make sure your family or carer is aware of the need to follow the advice below should sedation become a problem.

If you are awake and alert:	If you are mildly sedated and easy to rouse:	If you are frequently asleep, drowsy on waking but will follow commands or unable to remain awake:	If you are difficult to rouse or unarousable, or unable to, or finding it difficult, to follow commands:
No action is required	No action is required	Contact your GP or phone 111 for advice	Phone 999

Contact your GP or phone 111 if you are concerned about any side effect.

Can I have an alcoholic drink?

Do not drink any alcohol as this could increase drowsiness.

Will I become addicted?

If you are taking an opioid medicine to relieve pain, and for a short time only, it is very unlikely that you will become addicted. It is important that you do not take more than you need. You should not need to take for longer than a week. If you are still in severe pain after a week, you should see your doctor.

How should I store opioid medicines at home?

- Keep in the original container and store at room temperature in a dry place.
- Keep out of the sight and reach of children.
- Never give your medicine to anyone else.

What should I do with any unused opioid medicine?

- Return any unused medicine to your community pharmacist for safe disposal.
- Do not flush unused medicines down the toilet or throw them away.

About this leaflet:

Author: Christine Dodd, Medication Safety Pharmacist & Anne Gilbert
Written: May 2020
Approved: May 2020
Amended: June 2020
Review date: May 2023
Edition: 2

If you have feedback regarding the accuracy of the information contained in this leaflet, or if you would like a list of references used to develop this leaflet, please email pals@dchft.nhs.uk



INTEGRITY | RESPECT | TEAMWORK | EXCELLENCE

© 2020 Dorset County Hospital NHS Foundation Trust
Williams Avenue, Dorchester, Dorset DT1 2JY